

# 2010年 英文誌掲載論文

Circulating levels of 8 cytokines and marine n-3 fatty acids and indices of obesity in Japanese, white, and Japanese American middle-aged men.	Sekikawa A, Kadowaki T, Curb JD, Evans RW, Maegawa H, Abbott RD, Sutton-Tyrrell K, Okamura T, Shin C, Edmundowicz D, Kadota A, Choo J, El-Saad A, Ueshima H, Kuller LH; ERA JUMP Study group.	J Interferon Cytokine Res.
Effects of hyperglycemia on oxidative stress and antioxidant potential in patients with type 2 diabetes	Ikebuchi M, Nishio Y, Maegawa H, Kashiwagi A	Diabetol Int.
Laparoscopic sleeve gastrectomy leads to rapid improvement of glucose tolerance and insulin secretion with enhanced glucagon-like peptide (GLP-1) secretion	Ugi S, Yamamoto H, Kusunoki C, Kamo A, Ikeda K, Hashimoto T, Nishio Y, Kasama K, Tani T, Kashiwagi A, Maegawa H	Diabetol Int
Resveratrol improves oxidative stress and protects against diabetic nephropathy through normalization of Mn-SOD dysfunction in AMPK/SIRT1-independent pathway.	Kitada M, Kume S, Imaizumi N, Koya D.	Diabetes.
Activated microglia in the subthalamic nucleus in hyperglycaemic hemiballism: a case report.	Maeda K, Katayama Y, Sugimoto T, Somura M, Kajino Y, Ogasawara K, Yasuda H.	J Neurol Neurosurg Psychiatry.
Multiple cerebral aneurysms caused by HIV-associated vasculopathy.	Sanada M, Koizumi Y, Hodohara K, Kawai H.	Intern Med.
Amitriptyline and lorazepam improved catatonia and occipital hypoperfusion in a patient with DLB.	Maeda K, Ogawa N.	Intern Med.
Effects of blood pressure and the renin-angiotensin system on platelet activation in type 2 diabetes	Takashi Uzu, Masayoshi Sakaguchi, Atsuko Tsuda, Aya Kadota, Yukiyo Yokomaku, Shinji Kume, Masami Kanasaki, Keiji Isshiki, Shin-ichi Araki, Toshiro Sugimoto, Hiroshi Maegawa, Atsunori Kashiwagi	Journal of Diabetes Investigation
Presence and functional role of the rapidly activating delayed rectifier K(+) current in left and right atria of adult mice.	Nakamura H, Ding WG, Sanada M, Maeda K, Kawai H, Maegawa H, Matsuura H.	Eur J Pharmacol.
Low concentration of 4-hydroxy hexenal increases heme oxygenase-1 expression through activation of Nrf2 and antioxidative activity in vascular endothelial cells.	Ishikado A, Nishio Y, Morino K, Ugi S, Kondo H, Makino T, Kashiwagi A, Maegawa H.	Biochem Biophys Res Commun.
Prognostic value of normal stress myocardial perfusion imaging and ventricular function in Japanese asymptomatic patients with type 2 diabetes--a study based on the J-ACCESS-2 database.	Matsuo S, Nakajima K, Yamasaki Y, Kashiwagi A, Nishimura T.	Circ J.
Long-term effects of pioglitazone on carotid atherosclerosis in Japanese patients with type 2 diabetes without a recent history of macrovascular morbidity.	Yamasaki Y, Katakami N, Furukado S, Kitagawa K, Nagatsuka K, Kashiwagi A, Daida H, Kawamori R, Kaku K.	J Atheroscler Thromb.
Elevated serum levels of interleukin-18 in patients with overt diabetic nephropathy: effects of miglitol.	Uzu T, Yokoyama H, Itoh H, Koya D, Nakagawa A, Nishizawa M, Maegawa H, Yokomaku Y, Araki SI, Abiko A, Haneda M.	Clin Exp Nephrol.
Oleate and eicosapentaenoic acid attenuate palmitate-induced inflammation and apoptosis in renal proximal tubular cell.	Soumura M, Kume S, Isshiki K, Takeda N, Araki S, Tanaka Y, Sugimoto T, Chin-Kanasaki M, Nishio Y, Haneda M, Koya D, Kashiwagi A, Maegawa H, Uzu T.	Biochem Biophys Res Commun.
A genome-wide association study in the Japanese population identifies susceptibility loci for type 2 diabetes at UBE2E2 and C2CD4A-C2CD4B.	Yamauchi T, Hara K, Maeda S, Yasuda K, Takahashi A, Horikoshi M, Nakamura M, Fujita H, Grarup N, Cauchi S, Ng DP, Ma RC, Tsunoda T, Kubo M, Watada H, Maegawa H, Okada-Iwabu M, Iwabu M, Shojima N, Shin HD, Andersen G, Witte DR, Jorgensen T, Lauritzen T, Sandbak A, Hansen T, Ohshige T, Omori S, Saito I, Kaku K, Hirose H, So WY, Beury D, Chan JC, Park KS, Tai ES, Ito C, Tanaka Y, Kashiwagi A, Kawamori R, Kasuga M, Froguel P, Pedersen O, Kamatani N, Nakamura Y, Kadowaki T.	Nat Genet.
Monocyte chemoattractant protein-1 (MCP-1) gene polymorphism as a potential risk factor for diabetic retinopathy in Japanese patients with type 2 diabetes.	Katakami N, Matsuhisa M, Kaneto H, Matsuoka TA, Imamura K, Ishibashi F, Kanda T, Kawai K, Osonoi T, Kashiwagi A, Kawamori R, Shimomura I, Yamasaki Y.	Diabetes Res Clin Pract.
Replication study for the association between four Loci identified by a genome-wide association study on European American subjects with type 1 diabetes and susceptibility to diabetic nephropathy in Japanese subjects with type 2 diabetes.	Maeda S, Araki S, Babazono T, Toyoda M, Umezono T, Kawai K, Imanishi M, Uzu T, Watada H, Suzuki D, Kashiwagi A, Iwamoto Y, Kaku K, Kawamori R, Nakamura Y.	Diabetes.
Association between urinary type IV collagen level and deterioration of renal function in type 2 diabetic patients without overt proteinuria.	Araki S, Haneda M, Koya D, Isshiki K, Kume S, Sugimoto T, Kawai H, Nishio Y, Kashiwagi A, Uzu T, Maegawa H.	Diabetes Care.
Effects of a fish-based diet on the serum adiponectin concentration in young, non-obese, healthy Japanese subjects.	Kondo K, Morino K, Nishio Y, Kondo M, Fuke T, Ugi S, Iwakawa H, Kashiwagi A, Maegawa H.	J Atheroscler Thromb.
Transcription factor AP-2beta inhibits expression and secretion of leptin, an insulin-sensitizing hormone, in 3T3-L1 adipocytes.	Fuke T, Yoshizaki T, Kondo M, Morino K, Obata T, Ugi S, Nishio Y, Maeda S, Kashiwagi A, Maegawa H.	Int J Obes (Lond).