

# 2015年 英文誌掲載論文

Effect of aging on muscle mitochondrial substrate utilization in humans.	Petersen KF, Morino K, Alves TC, Kibbey RG, Dufour S, Sono S, Yoo PS, Cline GW, Shulman GI.	Proc Natl Acad Sci U S A.
Serum level of LOX-1 ligand containing ApoB is associated with increased carotid intima-media thickness in Japanese community-dwelling men, especially those with hypercholesterolemia LOX-1 ligand and IMT in Japanese.	Okamura T, Miura K, Sawamura T, Kadota A, Hisamatsu T, Fujiyoshi A, Miyamatsu N, Takashima N, Miyagawa N, Kadokawa T, Ohkubo T, Murakami Y, Nakamura Y, Ueshima H; SESSA Research Group	J Clin Lipidol
Association between Pulse Wave Velocity and Coronary Artery Calcification in Japanese men.	Torii S, Arima H, Ohkubo T, Fujiyoshi A, Kadota A, Takashima N, Kadokawa S, Hisamatsu T, Saito Y, Miyagawa N, Zaid M, Murakami Y, Abbott RD, Horie M, Miura K, Ueshima H; SESSA Research Group..	J Atheroscler Thromb
Renoprotective effect of DPP-4 inhibitors against free fatty acid-bound albumin-induced renal proximal tubular cell injury.	Tanaka Y, Kume S, Chin-Kanasaki M, Araki H, Araki S, Ugi S, Sugaya T, Uzu T, Maegawa H.	Biochem Biophys Res Commun.
Genome-wide association studies in the Japanese population identify seven novel loci for type 2 diabetes.	Imamura M, Takahashi A, Yamauchi T, Hara K, Yasuda K, Grarup N, Zhao W, Wang X, Huerta-Chagoya A, Hu C, Moon S, Long J, Kwak SH, Rasheed A, Saxena R, Ma RC, Okada Y, Iwata M, Hosoe J, Shojima N, Iwasaki M, Fujita H, Suzuki K, Danesh J, Jørgensen T, Jørgensen ME, Witte DR, Brandslund I, Christensen C, Hansen T, Mercader JM, Flannick J, Moreno-Macías H, Burtt NP, Zhang R, Kim YJ, Zheng W, Singh JR, Tam CH, Hirose H, Maegawa H, Ito C, Kaku K, Watada H, Tanaka Y, Tobe K, Kawamori R, Kubo M, Cho YS, Chan JC, Sanghera D, Frossard P, Park KS, Shu XO, Kim BJ, Florez JC, Tusié-Luna T, Jia W, Tai ES, Pedersen O, Saleheen D, Maeda S, Kadowaki T.	Nat Commun.
Impaired podocyte autophagy exacerbates proteinuria in diabetic nephropathy.	Tagawa A, Yasuda M, Kume S, Yamahara K, Nakazawa J, Chin-Kanasaki M, Araki H, Araki S, Koya D, Asanuma K, Kim EH, Haneda M, Kajiwara N, Hayashi K, Ohashi H, Ugi S, Maegawa H, Uzu T.	Diabetes.
Mammalian autophagy is essential for hepatic and renal ketogenesis during starvation.	Takagi A, Kume S, Kondo M, Nakazawa J, Chin-Kanasaki M, Araki H, Araki S, Koya D, Haneda M, Chano T, Matsusaka T, Nagao K, Adachi Y, Chan L, Maegawa H, Uzu T	Sci Rep.
Lipoprotein-associated phospholipase A2 is related to risk of subclinical atherosclerosis but is not supported by Mendelian randomization analysis in a general Japanese population.	Ueshima H, Kadowaki T, Hisamatsu T, Fujiyoshi A, Miura K, Ohkubo T, Sekikawa A, Kadota A, Kadowaki S, Nakamura Y, Miyagawa N, Okamura T, Kita Y, Takashima N, Kashiwagi A, Maegawa H, Horie M, Yamamoto T, Kimura T, Kita T; ACCESS and SESSA Research Groups.	Atherosclerosis.
Renal sinus fat volume on computed tomography in middle-aged patients at risk for cardiovascular disease and its association with coronary artery calcification.	Murakami Y, Nagatani Y, Takahashi M, Ikeda M, Miyazawa I, Morino K, Ohkubo T, Maegawa H, Nitta N, Sakai H, Nota H, Ushio N, Murata K.	Atherosclerosis
Emerging role of podocyte autophagy in the progression of diabetic nephropathy.	Yasuda-Yamahara M, Kume S, Tagawa A, Maegawa H, Uzu T.	Autophagy
MicroRNA-494 plays a role in fiber type-specific skeletal myogenesis in human induced pluripotent stem cells.	Iwasaki H, Imamura T, Morino K, Shimosato T, Tawa M, Ugi S, Sakurai H, Maegawa H, Okamura T.	Biochem Biophys Res Commun.
1-Methylnicotinamide ameliorates lipotoxicity-induced oxidative stress and cell death in kidney proximal tubular cells.	Tanaka Y, Kume S, Araki H, Nakazawa J, Chin-Kanasaki M, Araki S, Nakagawa F, Koya D, Haneda M, Maegawa H, Uzu T.	Free Radic Biol Med.
Urinary Potassium Excretion and Renal and Cardiovascular Complications in Patients with Type 2 Diabetes and Normal Renal Function.	Araki S, Haneda M, Koya D, Kondo K, Tanaka S, Arima H, Kume S, Nakazawa J, Chin-Kanasaki M, Ugi S, Kawai H, Araki H, Uzu T, Maegawa	Clin J Am Soc Nephrol.
Pioglitazone treatment and cardiovascular event and death in subjects with type 2 diabetes without established cardiovascular disease (JDDM 36).	Yokoyama H, Araki S, Kawai K, Hirao K, Oishi M, Sugimoto K, Sone H, Maegawa H, Kashiwagi A; Japan Diabetes Clinical Data Management Study Group.	Diabetes Res Clin Pract.
Duality of n-3 Polyunsaturated Fatty Acids on Mcp-1 Expression in Vascular Smooth Muscle: A Potential Role of 4-Hydroxy Hexenal.	Nagayama K, Morino K, Sekine O, Nakagawa F, Ishikado A, Iwasaki H, Okada T, Tawa M, Sato D, Imamura T, Nishio Y, Ugi S, Kashiwagi A, Okamura T, Maegawa H.	Nutrients.
Effect of nutritional counseling and long term isomaltulose based liquid formula (MHN-01) intake on metabolic syndrome.	Takeda E, Yamanaka-Okumura H, Taketani Y, Inagaki N, Hosokawa M, Shide K, Maegawa H, Kondo K, Kawasaki E, Shinozaki S, Fujinaka Y, Matsubara T, Katayama T, Sasaki H, Kawashima A, Aonuma H.	J Clin Biochem Nutr.
Focal Segmental Glomerular Sclerosis Ameliorated by Long-term Hemodialysis Therapy with Low-density Lipoprotein Apheresis.	Araki H, Ono S, Nishizawa Y, Deji N, Nakazawa J, Morita Y, Kume S, Chin-Kanasaki M, Isshiki K, Araki S, Arimura T, Maegawa H, Uzu T.	Intern Med
Lamp-2 deficiency prevents high-fat diet-induced obese diabetes via enhancing energy expenditure.	Yasuda-Yamahara M, Kume S, Yamahara K, Nakazawa J, Chin-Kanasaki M, Araki H, Araki S, Koya D, Haneda M, Ugi S, Maegawa H, Uzu T.	Biochem Biophys Res Commun.
Population Pharmacokinetics and Therapeutic Efficacy of Febuxostat in Patients with Severe Renal Impairment.	Hira D, Chisaki Y, Noda S, Araki H, Uzu T, Maegawa H, Yano Y, Morita SY, Terada T.	Pharmacology.
Predictors for Mild and Severe Hypoglycemia in Insulin-Treated Japanese Diabetic Patients.	Sonoda N, Morimoto A, Ugi S, Morino K, Sekine O, Nemoto K, Godai K, Maegawa H, Miyamatsu N, Matsuba R, Sakai K, Imamura M, Tanaka Y, Iwata M, Hirose H, Kaku K, Maegawa H, Watada H, Tobe K, Kashiwagi A, Kawamori R, Maeda S.	PLoS One.
Replication Study in a Japanese Population to Evaluate the Association between 10 SNP Loci, Identified in European Genome-Wide Association Studies, and Type 2 Diabetes.	Okano J, Kojima H, Katagi M, Nakae Y, Terashima T, Nakagawa T, Kurakane T, Okamoto N, Morohashi K, Maegawa H, Udagawa J.	PLoS One.
Epidermis-dermis junction as a novel location for bone marrow-derived cells to reside in response to ionizing radiation.		Biochem Biophys Res Commun.